

# Conformational isomerism in 3,5,8-trioxabicyclo[5.1.0]octane and its diastereomeric 4-methyl derivatives. A combined IR, X-ray and ab initio study

A.E. Klimovitskii<sup>a</sup>, E.E. Zvereva<sup>b</sup>, G.A. Chmutova<sup>a</sup>, A.B. Dobrynin<sup>b</sup>, V.Yu. Fedorenko<sup>a</sup>, Yu.G. Shtyrin<sup>a</sup>, I.A. Litvinov<sup>b</sup>, T.V. Bulygina<sup>a</sup>, E.N. Klimovitskii<sup>a,\*</sup>

<sup>a</sup> A.M. Butlerov Chemical Institute, Kazan State University, Kazan 420111, Box 115, Russian Federation

<sup>b</sup> A.E. Arbusov Institute of Organic and Physical Chemistry, Russian Academy of Sciences, Arbuzov Str. 8, Kazan 420088, Russian Federation

Received 27 March 2006; received in revised form 25 May 2006; accepted 30 May 2006

Available online 4 August 2006

## Abstract


IR spectroscopy and ab initio calculations have been applied to the title compounds. Bicyclic acetals were found to exist as mixture of chair and twist-boat conformations, a parent epoxide being in ternary equilibrium. X-ray data on 4-methyl (exo) derivative display a twist-boat form.

© 2006 Elsevier B.V. All rights reserved.

**Keywords:** Seven-membered bicyclic acetals; IR spectroscopy; X-ray crystallography; ab initio calculations

## 1. Introduction

Investigations of 1,3-dioxanes with substituent at C<sup>5</sup> (polar especially) show a number of interesting trends [1]. 3,5-Dioxabicyclo[5.1.0]octanes with exo- and endo chair-like conformations relate closely to above 1,3-dioxanes if one assumes that three-membered cycles in these molecules are equivalent to substituents at C<sup>5</sup> in the stereochemical sense (axial  $\equiv$  endo and equatorial  $\equiv$  exo). The noticeable feature of bicyclic acetals is the real propensity to exist in both chair conformations and a twist-boat form in addition [2–9]. It is worth noting that seven-membered acetals with a planar fragment have been inspected thoroughly and untraditional chair–twist-boat (C  $\rightleftharpoons$  TB) equilibrium has been clarified [10].



One can expect that bicyclic acetals with X = C, O, S, N may serve as suitable models for the investigation of polar and steric interactions between acetal moiety and three-membered rings. Encouraged by available stereochemical information implying both a broad range of

\* Corresponding author. Fax: +7 8432 9272 78.

E-mail address: [evgenii.klimovitskii@ksu.ru](mailto:evgenii.klimovitskii@ksu.ru) (E.N. Klimovitskii).